

KEEP OUT OF REACH OF CHILDREN

Blossom Boost

Liquid Fertiliser ideal for all Flower, Fruit and Vegetable Crops

2%N - 5%P - 10%K +TE <500 PPM (Cu,B,Fe,Na)

Important:

Spray tank must be cleaned and free from other chemical residuals.

Use cold water

Do not mix with lime sulphur

Compatibility: May be tank mixed with most commonly used agrochemicals. Please do jar test first.

Application Time and Rate

First Application: at 10 – 15% of flowering at the rate of 2.5L per ha mix with minimum 500L water per ha. Water can be increased up to 1000L/ha depends on crop type and application system.

Second Application: at 40 – 50% of flowering at the rate of 2.5L per ha mix with minimum 500L water per ha. Water can be increased up to 1000L/ha depends on crop type and application system.

Third Application: at 70 – 80% of flowering at the rate of 2.5L per ha mix with minimum 500L water per ha. Water can be increased up to 1000L/ha depends on crop type and application



Group Limited

Manufactured By: BSL

Call Free: 0800 77 55 11

Fax: 06 370 3527

Postal: P O Box 290. Masterton

www.mybsl.co.nz



Mixing Instructions

1. Put 1/3 of water in tank.
2. Add other chemicals, if any.
3. Fill tank with balance of water.
4. Add correct amount of fertiliser.
5. Agitate adequately to mix.
6. All analysis w/v.
7. Shake well before use.

Check compatibility with orthophosphate, high potash suspension, and chemical mixtures.