

KEEP OUT OF REACH OF CHILDREN

**Liquid Fertiliser
Manassium 12%w/v
Plus TE < 500 PPM (Cu,B,Fe,Na,Zn)**

MagMore

Ideal to soil application before planting

Magnesium (Mg) is a critical structural component of the chlorophyll molecule and is necessary for functioning of plant enzymes to produce carbohydrates, sugars and fats. It is used for fruit and nut formation and essential for germination of seeds. Deficient plants appear chlorotic, show yellowing between veins of older leaves; leaves may droop. Contains cat ions which improve moisture to the soils.

Soil Application: 10 - 50L/ha, mix with 1:20 (1 part of MagMore and 10 parts of water). Apply after grazing or bare land or pre planting.

Aerial Application: Ratio 1-2(10L per hectare mix with 20L of water).

Apply in Spring or Autumn, in cool weather.

Compatibility: May be tank mixed with most commonly used agrochemicals.

Withholding Period: 0 day.

Storage: Keep from freezing.

Directions For Use

After application, wash the solution off from spraying system. Agitation can be accomplished with a slow speed mechanical agitator, pump re-circulation, air, etc. Slurry is relatively easy to keep in suspension. It is also relatively easy to re-agitate if the material does settle. In more than 100L / 200L open lid barrel can be mixed with normal garden rake by agitating up and down.

Storage: can be easily stored in mild steel or poly tanks. Keep in a tightly closed container, stored in a cool, dry, ventilated area. Protect against physical damage. Isolate from incompatible substances.

Inhalation:

Remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Call a physician immediately.

Ingestion:

DO NOT INDUCE VOMITING. Give large quantities of water. Never give anything by mouth to an unconscious person. Call a physician immediately.

Skin Contact:

In case of contact, wipe off excess material from skin then immediately flush skin with plenty of water for at least 15 minutes. Remove contaminated clothing and shoes. Wash clothing before reuse. Call a physician immediately.

Eye Contact:

Immediately flush eyes with gentle but large stream of water for at least 15 minutes, lifting lower and upper eyelids occasionally. Call a physician immediately.

Mixing Instructions

1. Put 1/3 of water in tank.
2. Add other chemicals, if any.
3. Fill tank with balance of water.
4. Add correct amount of fertiliser.
5. Agitate adequately to mix.
6. All analysis w/v.
7. Shake well before use.

Check compatibility with orthophosphate, high potash suspension, and chemical mixtures.



**194 Bruntwood Rd, RD1. Cambridge.
P O Box 7071, Hamilton East. Hamilton.
0800 77 55 11 - Fax 0064 7 839 6365
www.mybsl.co.nz**

Distributor